How to give the right amount of ACETAMINOPHEN (also known as Tylenol) is different depending on which medicine you plan to give.

Dose: Give every 4 to 6 hours as needed for fever or pain. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Do NOT use with any other medicine containing acetaminophen.

Weight	Age	Infant's Acetaminophen (160 mg / 5 mL)	Children's Acetaminophen (160 mg / 5 mL)	Children's Acetaminophen Chewables (160 mg)	Children's Acetaminophen Dissolvable Packets (160 mg / powder pack)	Adult's Acetaminophen Tablets (325 mg)	Adult's Acetaminophen Tablets (500 mg)
6 to 11 pounds (3 to 5 kilograms)	0 to 3 months	1.25 mL					
12 to 17 pounds (about 5 to 7 kilograms)	4 to 11 months	2.5 mL					
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	3.75 mL					
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years		5 mL → 15 mL - 12.5 mL - 15 mL - 7.5 mL - 15	1 tablet			
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years		-15 mL- -10 mL- -5 mL- -7.5 mL	1 ½ tablets			
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years		10 mL → 15 mL − 12.5 mL − 15 mL − 7.5 mL − 10	2 tablets	2 packets	1 tablet	
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years		-15 mL- -10 mL- -5 mL- -7.5 mL-	2 ½ tablets	2 packets	1 tablet	
72 to 95 pounds (about 33 to 43 kilograms)	11 years		15 mL → 15 mL − 12.5 mL − 10 mL − − 15 mL − − 7.5 mL −	3 tablets	3 packets	1 ½ tablets	1 tablet
96 pounds or more (more than 43 kilograms)	12 years or older		$ \begin{array}{c} 10 \text{ mL} \rightarrow \begin{array}{c} 15 \text{ mL} - 12.5 \text{ mL} \\ 10 \text{ mL} - 7.5 \text{ mL} - \\ -5 \text{ mL} - 7.5 \text{ mL} \end{array} $ $ \begin{array}{c} and \\ 10 \text{ mL} \rightarrow \begin{array}{c} 15 \text{ mL} - 12.5 \text{ mL} \\ -5 \text{ mL} - 7.5 \text{ mL} - \\ -5 \text{ mL} - 7.5 \text{ mL} \end{array} $ $ \begin{array}{c} 10 \text{ mL} + 10 \text{ mL} = 20 \text{ mL} \end{array} $	4 tablets		2 tablets	1 tablet

How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.** Do **NOT** use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months				
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL	2.5 mL*		
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL [<u>***</u> *	4 mL*		
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL	5 mL* →	1 tablet	
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL	15stc - 12.5stc 10stc - 7.5stc 5stc - 7.5stc	1 ½ tablets	
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL	10 mL* → 15 mL - 12.5 mL 10 mL*	2 tablets	1 tablet
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years		-15sic12.5sic 10sic7.5sic7.5sic	2 ½ tablets	1 tablet
72 to 95 pounds (about 33 to 43 kilograms)	11 years		15 mL*	3 tablets	1 ½ tablets
96 pounds or more (44 kilograms or more)	12 years or older		10 mL + 10 mL = 20 mL* 10 mL* → 15 nL - 12.5 nL and 10 mL* → 15 nL - 12.5 nL 10 mL* → 15 nL - 12.5 nL - 15 nL - 12.5 nL - 15 nL - 12.5 nL - 15 nL - 12.5 nL	4 tablets	2 tablets

^{*}Note: This dosage is for **children's liquid ibuprofen** products given by dropper. See the previous column for correct dose-by-weight for **infant ibuprofen drops**, which are more concentrated.

How to give the right amount of DIPHENHYDRAMINE (also known as Benadryl) is different depending on which type of Diphenhydramine you plan to give.

Dose: Give every 6 hours if needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Do **NOT** use with any other medicine with diphenhydramine in it.

1	Be to a doe wat any other measure with apricing a anime in the									
Children's Liquid Diphenhydramine (12.5 mg / 5 mL)	Children's Diphenhydramine Chewable Tablets (12.5 mg)	Diphenhydramine Tablets (25 mg)	Diphenhydramine Capsules (25 mg)							
4 mL										
5 mL → -7.5 mL	1 tablet	½ tablet								
-10 mL -5 mL − 7.5 mL	1 ½ tablets	½ tablet								
10 mL → 10 mL - 7.5 mL -	2 tablets	1 tablet	1 capsule							
	4 tablets	2 tablets	2 capsules							
	Diphenhydramine (12.5 mg / 5 mL) 4 mL 3 mL -10 mL -7.5 mL -7.5 mL -7.5 mL 7.5 mL	Diphenhydramine (12.5 mg / 5 mL) 4 mL 5 mL 1 tablet 1 1/2 tablets 10 mL 7.5 mL 2 tablets 2 tablets	Diphenhydramine (12.5 mg / 5 mL) Chewable Tablets (12.5 mg) (25 mg) 4 mL 4 mL 1 tablet ½ tablet 5 mL 5 mL 1½ tablets ½ tablet 1 1½ tablets ½ tablet 1 tablet 2 tablets 1 tablet 1 tablet							

Do not give to children less than 2 years of age.

Do not give to children 2 to 6 years of age unless your doctor tells you to.